

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01K 67/027, 67/033, C12N 15/00	A2	(11) International Publication Number: WO 98/28971 (43) International Publication Date: 9 July 1998 (09.07.98)
(21) International Application Number: PCT/US97/23819 (22) International Filing Date: 31 December 1997 (31.12.97) (30) Priority Data: 08/778,645 3 January 1997 (03.01.97) US 60/080,222 3 January 1997 (03.01.97) US (71) Applicant (for all designated States except US): UNIVERSITY TECHNOLOGY CORPORATION [US/US]; Suite 250, 3101 Ins Avenue, Boulder, CO 80301 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): LINK, Christopher [US/US]; 1837 Spruce Street, Boulder, CO 80302 (US). (74) Agents: MONROY, Gladys, H. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: BETA-AMYLOID TOXICITY (57) Abstract <p>The present invention relates to methods and compositions for visualization of the toxic effects of transgenes <i>in vivo</i>. In particular, the present invention provides methods and compositions for the production and use of transgenic, including dually transgenic, <i>Caenorhabditis elegans</i> for visualization of the toxic effects of β-amyloid accumulation <i>in vivo</i>.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BETA-AMYLOID TOXICITY

FIELD OF THE INVENTION

The present invention is in the field of medicine and molecular biology. In particular, the invention provides transgenic non-human animals in which expression of a reporter gene is induced by a toxic transgene. The invention also provides methods and compositions for *in vivo* visualization of the toxicity associated with toxic proteins. In addition, the present invention provides methods and compositions for the production and use of transgenic non-human animals for screening candidate drugs to assess and treat toxicity.

BACKGROUND OF THE INVENTION

Investigation of the molecular mechanisms of various diseases has suggested that many pathologies are caused by expression of proteins that are immediately or gradually deleterious to the cells where they are expressed. Among these diseases are Alzheimer's disease, prion diseases, Huntington's disease, and amyotrophic lateral sclerosis. Several of these diseases are thought to result from aberrant folding of proteins, which results in the accumulation of toxic proteins or protein aggregates.

Alzheimer's disease is associated with the majority of dementia cases in the United States, with an estimated 2 million people afflicted with the disease, and a mortality rate of approximately 100,000 people per year (See, R.W.P. Cutler, "Degenerative and Hereditary Diseases," in *Medicine*, Scientific American, New York, (1988), pages 11 (IV):1-13; and R. Katzman (1986) *N. Engl. J. Med.* **314**:964). It has been estimated that the total cost for nursing home care alone of Alzheimer's patients exceeds \$13 billion/year (See, M.M. Heckler (1985) *Am. Psychol.* **40**:1240). According to the Centers for Disease Control (CDC), mortality due to Alzheimer's disease in the United States increased 10-fold between 1979 and 1987 (0.4 per 100,000 to 4.2 per 100,000) (See, "Reported death rate for Alzheimer's is up tenfold since 1979," (1990) *Clin. Psychiatr. News* **18**:21).

Patients suffering from Alzheimer's disease typically suffer progressive memory deficit, progressive decline in cognitive functions, anxiety, depression, visuospatial and speech deficits, delusions, personality changes, motor skill deterioration, loss of verbal

ability, and incontinence. Eventually, patients are completely incapacitated and disoriented, requiring total care. The course of Alzheimer's disease ranges from less than three years to over 20 years before death occurs. However, in typical cases, it progresses at a fairly constant rate, with an average duration of 6 to 10 years.

5 Alzheimer's disease is not the only cause of dementia. Indeed, there are over fifty recognized causes of dementia. As some causes of dementia are amenable to treatment, differential diagnosis of patients suffering dementia is particularly important. Because at least 20% of clinically diagnosed patients were found at autopsy to have had conditions other than Alzheimer's disease, the National Institute of Neurological and Communicative
10 Disorders and Stroke (NINCDS) and Alzheimer's Disease and Related Disorders Association (ADRDA) refined the clinical diagnostic criteria for Alzheimer's disease (See, M.A. Jenike, "Psychiatry," in *Medicine*, Scientific American, New York, [1991], pages 13 (V):1-5). Based on these criteria, the diagnosis of Alzheimer's disease may be "definite" (*i.e.*, requiring examination of brain tissue), "probable" (*i.e.*, patients have deficits in two or
15 more areas of cognition, insidious onset of disease, progressive worsening of memory and other cognitive functions, and normal consciousness levels), or "possible" (*i.e.*, patients meet the criteria for probable Alzheimer's disease, but exhibit variations in the disease course or have a systemic illness that is sufficient to cause dementia, but is not considered to be the cause of the dementia).

20 The numerous varieties of dementia, and variations in patient presentations, often make diagnosis problematic. Thus, the NINCDS/ADRDA criteria are very detailed, and necessitate the thorough examination of patients with suspected Alzheimer's disease. Currently, the only way to obtain a definite diagnosis is by post-mortem histological examination of brain tissue for the presence of senile plaques.

25 The histopathological lesions of Alzheimer's disease include neuritic or senile plaques, neurofibrillary degeneration, and granulovacuolar neuronal degeneration. The senile plaques usually contain a core of insoluble, amyloidic extracellular material (" β -amyloid") surrounded by a halo of neurofibrillary tangles and dystrophic neurons. The primary protein component of the amyloidic core of senile plaques is a 4.2 kd amyloid β
30 peptide (often referred to as " $A\beta$ "). A number of other proteins have also been identified

as components of senile plaques, including anti-chymotrypsin and apolipoprotein E. The major component of the neurofibrillary tangles is an abnormally phosphorylated microtubule-associated protein referred to as "tau." The β -amyloid protein in senile plaques is a small glycoprotein (*i.e.*, a 39-43 amino acid protein, derived from the larger amyloid precursor protein), that has been detected in non-neural tissues (*e.g.*, skin, subcutaneous tissue, and intestines) and blood vessels of Alzheimer's disease patients (*See*, Cutler, *supra*). Deposits of amyloid may be detected by their ability to bind specific dyes, such as Congo red or thioflavin S. Thus, it has been further hypothesized that detection of this protein may serve as a potential diagnostic aid in the assessment of patients suffering from dementia.

Alzheimer's disease usually arises spontaneously, although genetics play a role in development of the disease. For example, the histopathologic lesions of Alzheimer's disease have been regularly observed in the brains of older patients dying of Down's syndrome (*See*, Cutler, *supra*). It is known that the β -amyloid gene resides on chromosome 21. As Down's patients have an extra copy of this chromosome, there is an increased expression of this chromosome in these patients. It has been hypothesized that increased expression of the β -amyloid gene may regulate the formation of amyloid plaques ("senile plaques") in these patients. In addition, four loci were recently identified as playing a role in the genetic susceptibility of Alzheimer's disease (*See*, Pericak-Vance and Haines (1995) *Trends Genet.* 11:504).

Other potential risk factors for the development of Alzheimer's disease include environmental factors (*e.g.*, head trauma, smoking, and exposure to heavy metals), sociological factors (*e.g.*, depression and educational level), biological factors (*e.g.*, increasing age and hyperthyroidism), and a family history of Alzheimer's disease, Down's syndrome, or Parkinson's disease (*See*, Pericak-Vance and Haines, *supra*). Nonetheless, despite recent advances, the exact etiology and pathogenesis of Alzheimer's disease remain largely unknown.

In addition to the problems associated with diagnosing Alzheimer's disease, improvements are needed in the area of treatment. Many agents have been tested for their ability to treat the cognitive decline associated with Alzheimer's disease. For example,

various cholinergic enhancers (e.g., choline and lecithin) have been tested. Unfortunately, cholinergic precursors have been shown to be not useful, although some drugs that stimulate cholinergic transmission may be helpful in some patients. One example is physostigmine, a compound that prevents the synaptic breakdown of acetylcholine.

5 However, the overall clinical effect of this drug has not been as dramatic as initially hoped (See, M.A. Jenike, "Psychiatry," in *Medicine*, Scientific American, New York, (1991), pages 13 (V):1-5).

Other drugs, such as tetrahydroaminoacridine (THA or tacrine), a centrally acting anti-cholinesterase, have been tested. In a large multi-center trial of THA, liver enzyme
10 abnormalities were reported and the preliminary results indicated that, at least at low dosages, THA is not an effective treatment of Alzheimer's disease (Jenike, *supra*). Ergoloid mesylates (Hydergine) is an extremely safe compound, and remains the most commonly prescribed drug for patients with Alzheimer's disease. However, the overall effects of the drug are at best minimal.

15 The mechanisms of toxicity due to accumulation of β -peptide are currently the subject of much investigation, and no definitive causes of toxicity have yet been established. (See, Benzi and Moretti (1995) *Neurobiology of Aging*, 16:661-674). Because of this uncertainty, efforts to develop mechanism-based treatment regimens have not been possible. Compounds presently in use to treat Alzheimer's disease only serve to
20 alleviate the systemic effects associated with the disease.

In order to study disease mechanisms and genetic-based phenomena, animals in which a foreign gene has been inserted have been described by various researchers. International Patent Application WO 96/03034 describes insertion of retroviral vectors into fish, in order to produce fish with desirable traits or to study development. Various
25 transgenic animal models for Alzheimer's disease are described in International Patent Applications WO 93/14200, WO 93/02189, WO 94/12627, WO 94/23049, and European Patent Publication EP 653154. Typically these transgenic animals are mice or other mammals; however, β -amyloid peptide has been expressed in the nematode *Caenorhabditis elegans* (Link (1995) *Proc. Natl. Acad. Sci.* 92: 9368). The use of *C.*
30 *elegans* for investigation of mutant forms of the *C. elegans* genes *mec-4* and *deg-1*, which

cause neurodegeneration, is described in U.S. Patent No. 5,196,333. Drawbacks to mammalian animals as model systems are the relatively long generation time, which makes mammals less desirable for high-throughput screening of potential pharmaceuticals, and the difficulty in studying the molecular processes of interest without sacrificing the animal for cell and tissue analysis.

Reporter genes are genes that encode proteins or other compounds that can be detected by a variety of methods, and which "report" the occurrence of successful introduction and expression of gene sequences. β -galactosidase and luciferase are examples of such reporter genes. Recently, the gene for green fluorescent protein (GFP) of the jellyfish *Aequorea victoria* has been described in U.S. Patent No. 5,491,084. This gene can provide a method for indicating expression of a gene via fluorescent detection of GFP. The drawback to using GFP in mammalian systems is as described above, that is, the need to sacrifice the mammal in order to analyze the cells and tissues of interest. This drawback is especially severe in the context of large-scale screening of potential therapeutic compounds, and U.S. Patent No. 5,491,084 does not describe a system suitable for high-throughput screening of pharmaceuticals for activity against toxic proteins expressed in cells.

Thus, methods and animal systems are needed to screen drugs quickly and inexpensively for their effects on proteins and other substances associated with Alzheimer's disease. Convenient methods and animal systems for screening drugs for other neurodegenerative diseases, such as prion diseases, Huntington's disease, and amyotrophic lateral sclerosis, are also desirable.

SUMMARY OF THE INVENTION

The present invention provides methods and compositions useful for the production and use of dually transgenic animals, in particular *Caenorhabditis elegans*. In addition, in one embodiment, the present invention provides transparent animals that express a reporter gene inducible by a toxic transgene. In a preferred embodiment, the toxic transgene encodes β -peptide. Although it is not intended that the present invention be limited to any particular reporter, in a preferred embodiment, the reporter gene is green fluorescence

protein (GFP). It is contemplated that the effects or product of the reporter gene be observable (*i.e.*, the reporter gene is expressed). For example, in embodiments in which GFP is the reporter, its presence is detected using fluorescence microscopy. In addition, in these embodiments, the animals do not need to be sacrificed in order to observe the
5 expression of the reporter gene (*i.e.*, the detection may be accomplished on living animals). If other reporters are used, other detection methods may be necessary. For example, *lacZ* expression may be detected by exposing the tissues of the animal to the substrate for the gene (*i.e.*, β -galactoside), and observing for the presence of blue dye in the tissues. However, this method requires that the animal be sacrificed in order to observe the
10 expression of the reporter gene.

In one alternate embodiment, the genome of the transparent animal further contains a heterologous marker gene. In a preferred embodiment, the marker gene is *rol-6(su-1006)*. In the embodiments in which *rol-6(su-1006)*, expression of the marker gene is accomplished by observing the living animals for the rolling behavior that results due to
15 the expression of the abnormal collagen gene. In contrast, wild-type animals exhibit normal locomotion and do not exhibit this rolling behavior, as their collagen is normal.

In an alternative embodiment, the first transgene of the dually transgenic non-human animal comprises pCL25. In another alternative embodiment, the second transgene of the dually transgenic non-human animal comprises pCL12. In one alternate
20 embodiment, the genome of the transparent animal further contains a heterologous marker gene. In a preferred embodiment, the marker gene is *rol-6(su-1006)*. In the embodiments in which *rol-6(su-1006)*, expression of the marker gene is accomplished by observing the living animals for the rolling behavior that results due to the expression of the abnormal collagen gene. In contrast, wild-type animals exhibit normal locomotion and do not
25 exhibit this rolling behavior, as their collagen is normal.

In a preferred embodiment, the transparent animal is selected from the class Nematoda. Although it is not intended that the transparent animal of the present invention be limited to any specific animal, in a particularly preferred embodiment, the transparent animal is *Caenorhabditis elegans*.

In an alternate embodiment the cells of the transparent animal display toxicity resulting from the accumulation of β -peptide within the cells. In one preferred embodiment, the genome of the transparent animal comprises SEQ ID NO:5 and SEQ ID NO:8.

5 The present invention also provides methods for producing dually transgenic non-human animal comprising: providing: a first and second non-human animal; a first transgene comprising β -peptide; and a second transgene, comprising a reporter; introducing the first transgene into the genome of the first non-human animal to produce a first transgenic animal, and introducing the second transgene into the genome of the
10 second non-human animal to produce a second transgenic animal; and mating the first transgenic animal with the second transgenic animal to produce a dually transgenic animal, wherein the β -peptide and the reporter are expressed.

In one embodiment of the methods of the present invention the dually transgenic non-human animal is transparent. In a preferred embodiment, the transparent animal is a
15 nematode, while in a particularly preferred embodiment, the animal is *Caenorhabditis elegans*.

In an alternative embodiment, the first transgene of the dually transgenic non-human animal comprises pCL25. In another alternative embodiment, the second transgene of the dually transgenic non-human animal comprises pCL12. In one alternate
20 embodiment, the genome of the transparent animal further contains a heterologous marker gene. In a preferred embodiment, the marker gene is *rol-6(su-1006)*. In the embodiments in which *rol-6(su-1006)*, expression of the marker gene is accomplished by observing the living animals for the rolling behavior that results due to the expression of the abnormal collagen gene. In contrast, wild-type animals exhibit normal locomotion and do not
25 exhibit this rolling behavior, as their collagen is normal.

The present invention also comprises methods for testing compounds for anti-toxic effects, comprising: providing a dually transgenic non-human animal expressing a toxic transgene and a reporter; a composition comprising a test compound in a form suitable for administration such that the compound is bioavailable in the cells of the animal; and
30 administering the test compound to the non-human animal. The toxic transgene induces

expression of the reporter gene. In a preferred embodiment, the toxic transgene is β -peptide and the reporter gene is GFP. In one embodiment, the method further comprises the step of measuring a reduction or increase in the expression of the reporter by the dually transgenic non-human animal and thereby identifying a compound as therapeutic. In a particularly preferred embodiment of the methods, the compounds inactivate the β -peptide expressed by the dually transgenic animal.

In one embodiment of the methods for testing compounds for β -peptide toxicity, the dually transgenic non-human animal is transparent. In a preferred embodiment, the transparent animal is a nematode, while in a particularly preferred embodiment, the animal is *Caenorhabditis elegans*. In an alternative embodiment, the first transgene of the dually transgenic non-human animal comprises pCL25. In another alternative embodiment, the second transgene of the dually transgenic non-human animal comprises pCL12. In one alternate embodiment, the genome of the transparent animal further contains a heterologous marker gene. In a preferred embodiment, the marker gene is *rol-6(su-1006)*, which produces effects described herein.

In one alternative embodiment, one or more test compounds are tested for their ability to counter the toxic effects of transgene product. It is also contemplated that the test compounds will be tested for their ability to prevent the expression of the toxic transgene, for example, β -peptide.

In alternative embodiments, dually transgenic animals in which the expression of, or effects of transgene toxicity, are reduced or eliminated by the test compounds, are mated. The progeny of these matings are also then tested for the effects of test compounds on the expression of, or effects of toxicity. In yet another embodiment, the progeny of these matings are used in other assay systems for the identification of therapeutic compounds.

In yet another alternative embodiment, the dually transgenic animals of the present invention are used to identify methods suitable for the diagnostic testing of Alzheimer's disease or pathology due to other diseases. Thus, the dually transgenic animals of the present invention are used to develop assays suitable for use in humans or animal models of Alzheimer's disease.

DESCRIPTION OF THE FIGURES

Figure 1 is a diagram illustrating the construction of dual transgenic animals expressing β -amyloid peptide and an hsp/GFP reporter.

5 Figure 2 shows the DNA sequence (SEQ ID NO:5) and restriction map for pCL12.

Figure 3 is a graphic map of pCL12.

Figure 4 shows the DNA (SEQ ID NO:6), and amino acid sequence (SEQ ID NO:7) of pCL12 from nucleotide 1071 through 1253 (*i.e.*, the β -(1-42) nucleic acid and amino acid sequence).

10 Figure 5 shows the DNA sequence (SEQ ID NO:8) and restriction map for pGFP-TT.

Figure 6 is a graphic map of pGFP-TT.

Figure 7 shows the DNA sequence (SEQ ID NO:9) and restriction map for pCL25.

Figure 8 is a graphic map of pCL25.

15 Figure 9 shows the DNA sequence (SEQ ID NO:10) of *rol-6*.

Figure 10 shows the amino acid sequence (SEQ ID NO:11) of *rol-6*.

DESCRIPTION OF THE INVENTION

20 The present invention provides methods and compositions useful for the production and use of transgenic animals. The methods and transgenic animals of the invention also provide an efficient and effective system for screening drug effective in ameliorating the effects of toxic gene products.

25 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. *See, e.g.*, Sambrook, Fritsch, and Maniatis. MOLECULAR CLONING: A LABORATORY MANUAL, 2nd edition (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, (F.M. Ausubel et al. eds., 1987); the series METHODS IN ENZYMOLOGY (Academic Press, Inc.); PCR 2: A PRACTICAL APPROACH (M.J. McPherson, B.D. Hames and G.R. Taylor eds., 1995); ANIMAL CELL CULTURE (R.I.

30

Freshney, Ed., 1987), and ANTIBODIES: A LABORATORY MANUAL (Harlow et al. eds., 1987).

Animals

5 The methods and systems described herein can be practiced with any non-human animal. In a preferred embodiment, the animal used is transparent for at least part of its life cycle, for example, oocytes (*Xenopus* or others); larvae; pupae, fish (zebrafish or others); and nematodes, such as *C. elegans*. *C. elegans* is particularly preferred because it is transparent throughout its life and has been extensively studied as a model to determine
10 developmentally regulated gene expression, as well as pattern formation. The visibility of all of the animals' cells throughout their life cycles, has led to the complete determination of the cell lineages, and a detailed description of the morphogenic changes that occur during *C. elegans* development. Rapid methods for introduction of cloned DNA into the *C. elegans* germ line has provided means to study molecular function and expression *in*
15 *vivo*, as the effects of the gene expression are usually readily visible in the transparent worms.

Reporter Molecules

 The methods and compositions of the present invention make use of reporter genes
20 to monitor gene activity. Visible and quantifiable reporter genes are known and described in the art. Successfully used reporter molecules in gene fusion vectors in studies with *C. elegans* include the *E. coli lacZ* coding region (See e.g., A. Fire *et al.*, (1990) *Gene* **93**:189-198), and the *Aequora victoria gfp* coding region (See e.g., M. Chalfie *et al.*, (1994) *Science* **263**:802-805; D.C. Prasher *et al.*, (1992) *Gene* **111**:229-233; and Genbank
25 Accession #M62654), which produces an intrinsically fluorescent protein. Although various fusion expression vectors have been used and reported in the literature, problems have been encountered (See e.g., A. Fire, "Fire Lab Vector Kit"--June 1995). For example, with *lacZ*, there have been many reports in which it was not possible to correlate transgene expression patterns with physiological expectations, or only weak correlations have been
30 possible. Ectopic expression is often frequently seen with short promoters, and occurs

most prominently in the gut and pharynx of the affected animals. This may be due to weak promoter and/or enhancer signals in the vectors. Expression pattern deficits of *lacZ* fusions have been classified into three groups. In the first, transgenes are expressed in the correct tissue, but mosaic expression is observed (*i.e.*, only a subset of the cells stain during the detection methods). This has been observed even with integrated high-copy transgenes. In the second group, expression in a single tissue or cell population is not seen with the transgene. Finally, there has been the failure of transgene fusion constructs to show expression in the pre-12 cell embryo, or in any embryonic, larval, or adult germ line.

The *gfp* fusion vectors were developed as an alternative to the *lacZ* markers, but have been available for a much shorter time (*See e.g.*, Chalfie *et al.*, (1994), *supra*; U.S. Patent No. 5,491,084). Initial reports indicate that the fluorescence pattern appears to be more restricted than that exhibited by equivalent *lacZ* fusion. In addition, there are still problems associated with the germline expression of the transgenes and ectopic expression. Indeed, in some cases, the problems with ectopic expression have been exacerbated by these "improvements."

The present invention overcomes many of these problems described in the art. The transparent animals and dually transgenic animals described herein provide easily visible, stably expressed systems in which the toxic transgene and reporter gene are expressed similarly to endogenous chromosomal gene expression. These animals provide an excellent system for screening compounds having effects on the toxic transgene. The present invention, therefore, provides model systems for the study of human diseases and methods of identifying therapeutic compounds using these animal systems.

Inducible Promoters

As discussed above, the reporter genes are operably linked to an inducible promoter. The promoter is induced by the toxic transgene, for example by the gene product of the toxic transgene. It is also contemplated that the toxic transgene can act to induce the promoter indirectly, for example by disrupting other cellular proteins or functions. Suitable inducible promoters are available and can be readily determined by those skilled in the art. Non-limiting examples of promoters which are induced by

"stresses" include the metallothionein gene promoter (*mtl-1* or *mtl-2*; *e.g.*, Genbank Accession #M92910, #M11794, #X00504, and #X00953), and the *C. elegans* amyloid precursor protein (APP) homolog *apl-1* gene promoter (*See e.g.*, Daigle and Li, Proc. Natl. Acad. Sci., 90:12045-12049 [1993]). Other potential promoters include those from other
5 inducible heat-shock genes; at least one of the known *C. elegans hsp70* genes is strongly heat-inducible. It is also contemplated that promoters from genes known to be up-regulated under stress conditions in other systems (*e.g.*, superoxide dismutase, catalase, glutathione reductase, etc.) may also be useful.

In one embodiment, as described in the Examples below, a strong muscle-specific
10 promoter was used to express a potentially secretable form of the β -peptide, so as to generate significant extracellular levels of β -peptide. This was accomplished in order to mimic the situation that may exist in the human brain, and allow observations of cell-external neurotoxicity. The present inventor has also demonstrated that the promoter must be chosen so that enough β -peptide is produced to cause physiological effects, but not so
15 much to kill the animals.

The *unc-54*/ β -(1-42) minigene was constructed with a modified signal sequence that has been previously shown to allow secretion of a *her-1* protein product that is ectopically expressed in muscle cells (M.D. Perry *et al.*, (1993) *Genes Dev.*, 7:216-228). The development of *C. elegans* transgenic for expression of β -amyloid was previously
20 described by the inventor (Link, (1995) *Proc. Natl. Acad. Sci.*, 92:9368-9572).

However, in early experiments, it was observed that β -peptide deposits were not convincingly detected outside of the muscle cells, when tested with the antibodies described in Example 4, below. While an understanding of the mechanism is not necessary for the practice of the present invention, it is apparent that the majority of β -
25 peptide expressed by the transgenic worms is retained in the muscle cells and is responsible for the pathology observed in the muscle cells.

It is also contemplated that β -peptide expression will be directed to other tissues, through utilization of appropriate promoters. For example, it is contemplated that animals expressing β -peptide in the intestine may be particularly useful to analyze compounds such

as drugs for their effect on β -peptide, as these cells readily take up exogenous compounds administered orally.

Transgenes

5 The transgene element of the present invention can be any sequence which is able to induce the promoter operably linked to the reporter gene. As noted, the transgene is "toxic" in the sense that it disrupts cellular function in some way. Preferably, the transgene encodes a protein that is toxic to the host cells and/or organism in that it causes deleterious effects to the host, for instance interfering with the hosts ability to survive and/or grow. The toxic transgenes encode proteins that disrupt cellular function directly (e.g., the gene product is toxic) or indirectly (e.g., the sequence of the transgene disrupts cellular function by some mechanism other than its gene product). Toxic proteins are distinguishable from simple chemical toxins (e.g., heavy metals and the like) by their antigenicity and higher molecular weight.

15 In one embodiment, the toxic transgene comprises a gene encoding for an amyloidic protein, for example beta amyloid peptide, prion protein variants, transthyretin variants, gelsolin variants, cystatin variants, lysozyme variants and the like. In another embodiment, the transgene encodes a protein containing polyglutamine resulting from triplet-repeat expansion such as huntingtin (a protein that has been implicated in Huntington's Chorea), ataxin-1 or ataxin-2. Alternatively, proteins associated with inherent amyelotopic lateral sclerosis (ALS) for example, superoxide dismutase 1 variants and over-expressed neurofilament protein, can be used. It will be understood that the transgene can encode for an entire toxic protein or, alternatively, a functional (i.e. toxic) fragment.

Transgenic Animals

25 In one aspect, the present invention includes a dually transgenic non-human animal. In one embodiment, the development of these dually transgenic animals involves the production of two *C. elegans* lines with a single transgene in each line, designated as CL2005 and CL2070. *C. elegans* was chosen due to the observation that the toxic effect of

β -amyloid has been reported to occur in these animals, in addition to humans. Line CL2005 exhibited muscle-specific expression of human β peptide, while line CL2070 exhibited stress-inducible expression of GFP. GFP was incorporated into the present invention as it provides a marker (*i.e.*, "reporter") that is readily visible in living worms. Thus, dually transgenic animals both express β -peptide in their muscle cells and under appropriate conditions for observation (*i.e.*, fluorescence microscopy), exhibit green fluorescence in these cells.

Mating of these lines resulted in the production of dually transgenic animals, in which the presence of β -amyloid was easy to detect, due to the fact that the dually transgenic animals expressing β -amyloid glowed green. Thus, the present invention provides an easily detectable method for the specific expression of β -peptide in dually transgenic animals. This ease of detection provides great advantages for the development of methods to analyze the effects of β -amyloid *in vivo*. In addition, because the dually transgenic animals are not killed in order to detect the presence of β -amyloid, as would be required if other reporters (*e.g.*, *lacZ* or luciferase) were used, the animals may be useful for screening compounds for their effects on β -amyloid in an *in vivo* situation. Thus, the use of an reporter which is detectable *in vivo* provides significant advantages over currently available methods.

Other Advantages of Transgenic *C. elegans*

The present invention also includes methods of screening compounds for their ability to prevent or inhibit toxicity due to the toxic transgene. In these drug screening embodiments, the methods and animals described above can be used to identify potential therapies. For example, in assays to develop compounds to block the expression of β -amyloid or counter its toxic effects, transgenic worms (*e.g.*, green fluorescing worms with muscle-specific β -peptide expression), would be observed for their loss of fluorescence. In assays where the compound is effective in countering the expression of β -peptide or effects of β -peptide, the animals would no longer glow green. It is easy to visually screen and quantify (using commercially available equipment) for animals that have lost the ability to glow, and then use them to detail the interaction between the compound and the toxic

transgene. The present invention provides a fast and efficient screening system, for example by using commercially available equipment for assaying multiple compounds at once.

Thus, the transgenic animals of the present invention expressing a detectable reporter gene triggered by a toxic transgene provide animal models for human diseases. For instance, a reporter gene (such as GFP) induced by expression of β -peptide provides an animal model for human Alzheimer's disease. It is also not intended that the present invention be limited to animals expressing β -peptide in conjunction with GFP. It is contemplated that expression of other proteins associated with dementia and/or Alzheimer's or other diseases would also find use in conjunction with GFP in *C. elegans* as well. For example, it is contemplated that proteins (including mutated versions of proteins) such as the presenilins associated with Alzheimer's disease (See e.g., J. Marx (1996) *Science* 274:1838-1840) will be used in conjunction with GFP and full-length APP in dually transgenic *C. elegans*. Other toxic proteins which provide animal models for other human diseases are described herein.

These transgenic animals also provide a means to screen compounds for their ability to decrease or eliminate the toxicity. Examples of compounds suitable for testing using the transgenic animals of the present invention include, but are not limited to, Congo Red, tumor necrosis factor (TNF), estrogen, tacrine (9-amino-1,2,3,4-tetrahydroacridine), dihydroepiandrosterone (DHEA), compounds that inhibit ApoE4, and others. Compounds to be tested for anti-toxic activity are administered to the same number of dually transgenic animals (e.g., commercially available from suppliers such as Sigma. Compounds to be tested for anti-toxic activity are administered to the same number of dually transgenic animals (e.g., generated using the methods described in the Examples, below) from the control group and the treatment group, and the presence or absence of reporter gene used as a measure of efficacy.

The compounds being tested can be administered using any suitable route (e.g., oral, parenteral, controlled-release transdermal methods, and implants, etc.). In one preferred route, the compounds to be tested are suspended in the growth media provided to the worms. Generally speaking, the route of administration will depend on the stability of the compound, the susceptibility of the compound to "first pass" metabolism, the

concentration needed to achieve a therapeutic effect, and the like. Following initial screening, a compound that appears promising (*i.e.*, which increases the number of worms which display reduced β -peptide toxicity) is further evaluated by administering various concentrations of the compound to additional transgenic animals in order to determine an approximate therapeutic dosing range.

Another screening method involves the crossing of the transgenic worms of the present invention with other transgenic worms. The animals are observed after treatment, in the presence and absence of the test compound(s), with the effects on the toxic transgene being gauged either by crude survival or the presence/absence of the reporter gene. It is also contemplated that the methods of the present invention be modified so as to provide means to analyze disease-related proteins believed to have dominant toxic effects, by substituting appropriate sequences for the β -peptide sequences used as described in the Examples. For example, proteins such as transthyretin (known to be associated with familial amyloid polyneuropathy; *See e.g.*, Christmason *et al.*, (1991) *FEBS* **281**:177-180; Genbank Accession #D00096), and variant superoxide dismutase (known to be associated with familial amyotrophic lateral sclerosis [Lou Gehrig's disease]), prion proteins, A4 amyloid protein (*See e.g.*, Ponte *et al.*, (1988) *Nature* **331**:525-527; Salbaum *et al.*, U.S. Patent No. 5,151,508, herein incorporated by reference), APP (*See e.g.*, Kitaguchi *et al.*, (1988) *Nature* **331**:530-532; Sata *et al.*, EP Appln. 94117512.7; Scott *et al.*, WO 9412627; Wadsworth *et al.*, WO 9314200; Gearhart *et al.*, WO 9423049; and Neve *et al.*, WO 9302189, all of which are herein incorporated by reference), other amyloidic proteins (*e.g.*, variant lysozymes and amylin peptide), and other proteins associated with neuronal degeneration (*See e.g.*, U.S. Patent No. 5,196,333), will be used in the methods of the present invention by substituting the appropriate nucleic acid sequences encoding the protein of interest for the β -peptide described in Example 1. It is also contemplated that animals transgenic for dual proteins (*e.g.*, β -peptide in combination with another protein, such as transthyretin) will be used. Thus, the present invention provides the means to analyze the effects of numerous genes and proteins *in vivo*.

Furthermore, the transgenic animals of the present invention provide distinct advantages over other transgenic animals currently used to analyze diseases such as

Alzheimer's disease. Due to the short gestation period of *C. elegans*, transgenic animals can be produced much more rapidly than when mammals, such as mice are used. For example, transgenic mice overexpressing a 695 amino acid isoform of β -amyloid precursor did not show learning and memory impairment until they were 9-10 months of age (Hsiao *et al.*, Science 274:99-102 [1996]). In contrast, the transgenic animals of the present invention express β -peptide and may be manipulated at a very early age, even in embryonic stages.

In addition, because the transgenic animals are easy to select based on the presence of the green fluorescence, invasive procedures such as surgery, necessary to analyze the effects of the toxic transgene in other animals (e.g., mice) are avoided.

Finally, the apparent toxicity of β -peptide in transgenic animals was found to be temperature-dependent. Transgenic animals maintained at 25.5°C were significantly more sick than those maintained at 16°C. The animals maintained at the higher temperature became paralyzed more quickly, failed to eat or grow, failed to show normal egg-laying, and many died before reaching adulthood. However, wild-type animals grow well at either of these temperatures. The CL2005 parental line was temperature-sensitive for viability, as it was not possible to propagate this line at the elevated temperature. In the CL2070 parental line, 25.5°C maintenance was not sufficient to induce the hsp/GFP construct. The hsp/GFP response was similarly temperature-dependent. It was difficult to detect GFP when the dual transgenic animals were raised at 16°C, but GFP induction was dramatic when the animals were raised at 25.5°C.

These temperature dependency observations were exploited by the propagation of animals at 16°C, and then upshifting them to 25.5°C, when GFP induction was desired. This effect also has potential benefits in the analysis of compounds, as the animals can be pre-incubated in the presence of drugs or other compounds at the lower temperature before the upshift to the higher temperature, in order to ensure that the presumed protective effect of the drug was in place, prior to the strong induction of β -peptide toxicity. Importantly, these assays may be completed within one day and are suitable for rapid methods (e.g., the use of a microtiter format and a plate fluorimeter), so that literally thousands of compounds may be tested simultaneously.

In sum, the present invention provides methods and compositions useful as animal models for disease, as well as providing methods and compositions for disease therapy and prevention. The animal testing may be supplemented and confirmed by testing on human subjects. However, the transgenic animals of the present invention allow the testing of a large number of compounds, both various methods, including those known in the art.

DEFINITIONS

To facilitate understanding of the invention, a number of terms are defined below.

The term "non-human animal" includes vertebrates such as rodents, arthropods, insects (*e.g.*, Diptera), fish (*e.g.*, zebrafish), non-human primates, ovines, bovines, ruminants, lagomorphs, porcines, caprines, equines, canines, felines, aves, etc. The non-human animal can be used at any stage in its development, for example oocyte, fetal, larval, pupal stages or the like. Preferred non-human animals are those that are transparent, such as certain nematodes, oocytes, larvae and fish.

As used herein, the term "transparent" is used in references to animals through which light will be transmitted. However, it is not intended that the amount of light transmittance be limited to any particular amount. For example, an animal is transparent so long as at least some light may be transmitted through its body. Non-limiting examples of transparent animals are larval stages of some animals (*e.g.*, flies) and oocytes (*e.g.*, *Xenopus* oocytes). Particularly preferred non-human animals are selected from the nematodes (*i.e.*, any animal in the Class Nematoda), most preferably *Caenorhabditis elegans*. However, it is contemplated that other transparent animals, such as zebrafish will be useful in the present invention. For example, transgenic zebrafish have been produced (See *e.g.*, Lin *et al.*, PCT Publ. WO9603034; incorporated herein by reference).

The "non-human animals having a genetically engineered genotype" of the invention are preferably produced by experimental manipulation of the genome of the germline of the non-human animal. These genetically engineered non-human animals may be produced by several methods including the introduction of a "transgene" comprising nucleic acid (usually DNA) into an embryonal target cell or integration into a chromosome of the somatic and/or germ line cells of a non-human animal by way of human

intervention, such as by the methods described herein. The process by which a DNA molecule becomes stably incorporated into another genome is referred to as "stable integration." Non-human animals which contain a transgene are referred to as "transgenic non-human animals". A transgenic animal is an animal whose genome has been altered by the introduction of one or more transgenes.

The term "transgene" as used herein refers to a foreign gene that is placed into an organism by introducing the foreign gene into gonadal cells, embryonic cells, newly fertilized eggs, or early embryos. The term "foreign gene" refers to any nucleic acid (*e.g.*, gene sequence) which is introduced into the genome of an animal by experimental manipulations and may include gene sequences found in that animal so long as the introduced gene contains some modification relative to the naturally-occurring gene. A preferred foreign gene is the β -amyloid gene (*e.g.*, β -peptide), or fragments thereof. A "toxic transgene" refers to a foreign gene which disrupts cellular function in some way. For example, the toxic transgene may produce a gene product (*i.e.*, protein) which is toxic to the cell or organism. Alternatively, the transgene may disrupt other cellular proteins, or act as a DNA binding site without producing a protein.

As used herein, the term "toxicity" refers to the production of toxic effects by any compound or substance. For example, while it is not necessary to the understanding and use of the present invention, β -peptide toxicity may arise due to the accumulation of β -peptide molecules. There are numerous manifestations of toxicity that may occur. For example, the toxicity associated with β -peptide may be manifested as negative effects on muscle cells of dually transgenic animals.

The terms "promoter element," "promoter," or "promoter sequence" as used herein, refer to a DNA sequence that is located at the 5' end of (*i.e.*, precedes) a gene in a DNA polymer and provides a site for initiation of the transcription of the gene into mRNA. An "inducible promoter" is a promoter that is triggered by certain signals within the cell, for instance binding of a transcription factor, stress, heat or the like.

The term "reporter gene" as used herein refers to genes that encode proteins or other compounds that can be detected by a variety of methods. These genes "report" the occurrence of successful introduction and expression of sequences such as transgenes.

Non-limiting examples of reporter genes include antibiotic resistance genes, genes encoding enzymes and genes encoding other detectable proteins. Expression of the reporter gene is detected using methods known in the art. In a preferred embodiment, the reporter gene is a GFP gene, although *lacZ* β -galactosidase gene, or any other reporter system may be used to detect the successful production of transgenic animals. In particularly preferred embodiments, the reporter is a compound or protein which may be present or expressed within living animals. That is, it is not necessary to sacrifice the animal in order to detect the presence of the reporter. The type of the reporter gene system used is not critical to the invention, and it is contemplated that any system suitable for use with the transgenic animals of the present invention will be used.

As used herein, the terms "dual transgenic" and "dually transgenic" refer to animals or cells in which more than one transgene have been introduced. For example, the term is used in reference to cells which contain the sequences encoding β -peptide, and sequences encoding a reporter (*e.g.*, GFP). However, it is not intended that the number of transgenes in the dually transgenic animals of the present invention be limited to two. For example, the transgenic animals may also contain another one or more marker genes (*e.g.*, *rol-6* sequences), in addition to β -peptide and reporter sequences. In the embodiments in which *rol-6(su-1006)*, expression of the marker gene is accomplished by observing the living animals for the rolling behavior that results due to the expression of the abnormal collagen gene. In contrast, wild-type animals exhibit normal locomotion and do not exhibit this rolling behavior, as their collagen is normal.

The transgenic animals of the present invention are preferentially generated by introduction of the targeting vectors into gonad cells. Transgenes can be efficiently introduced into the cells by DNA transfection using a variety of methods known to the art, including electroporation, calcium phosphate co-precipitation, protoplast or spheroplast fusion, lipofection, and DEAE-dextran-mediated transfection. Transgenes may also be introduced into cells by retrovirus-mediated transduction or by micro-injection. In one preferred embodiment, the transgenes are injected into gonads of *C. elegans* as described by Mello *et al.* (1991) *EMBO J.* 10:3959-3970. Alternative methods for the generation of transgenic animals containing an altered gene are known to the art. For example,

embryonal cells at various developmental stages can be used to introduce transgenes for the production of transgenic animals. Different methods are used depending on the stage of development of the embryonal cell.

5 A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting related sequences include oligolabeling, nick translation, and end-labeling or PCR amplification using a labeled nucleotide. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles and the like. The term may also be used in reference to 10 proteins. For example, a variety of protocols which employ polyclonal or monoclonal antibodies specific for the β -peptide protein product are known in the art (*See, the Examples*). These antibodies can be used as markers for the expression of proteins. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). 15

The term "gene" refers to a DNA sequence that comprises coding sequences necessary for the production of a polypeptide or precursor. In some instances, a gene can also include control sequences. The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired enzymatic activity is retained.)

The term "gene of interest" refers to any gene, the manipulation of which may be deemed desirable for any reason, by one of ordinary skill in the art.

The term "wild-type" refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that naturally-occurring mutants can be isolated: these are

tified by the fact that they have altered characteristics when compared to the wild-type or gene product.

The terms "targeting vector" or "targeting construct" refer to oligonucleotide sequences comprising a gene of interest flanked on either side by regulatory sequences. Typically, the targeting vector is capable of homologous recombination such that the gene of interest is integrated by recombination.

As used herein, the terms "vector" and "vehicle" are used interchangeably in reference to nucleic acid molecules that transfer DNA segment(s) from one cell to another.

The terms "expression vector" or "expression cassette" as used herein, refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a host organism. Nucleic acid sequences necessary for expression in prokaryotes include a promoter, an operator (optional), and a ribosome binding site, often along with other regulatory sequences. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

The terms "in operable combination," "in operable order," and "operably linked" as used herein, refer to the linkage of nucleic acid sequences in such a manner that a nucleic acid molecule capable of directing the transcription of a given gene and/or the synthesis of a protein molecule is produced. The term also refers to the linkage of amino acid sequences in such a manner so that a functional protein is produced.

The term "tissue-type specific" as it applies to a promoter, refers to a promoter that is capable of selectively directing expression of a gene in a specific tissue. Similarly, the term "tissue-specific promoter" in the method of the present invention does not require absolute tissue specificity. In general, the requisite specificity is found where a plurality (or, typically, a majority) of cells in one tissue type express a gene of interest, while in other tissue types, e.g., greater than 80%, and preferably greater than 90%, and more preferably greater than 95%, of the cells in other tissue types do not. In one embodiment of the invention, the strong muscle promoter *unc54* was used (See, Example 1).

As used herein the term "portion" when in reference to a gene refers to fragments of a gene. Fragments may range in size from a few nucleotides to the entire gene.

sequence minus one nucleotide. Thus, "an oligonucleotide comprising at least a portion of a gene" may comprise fragments of the gene or the entire gene.

As used herein, the term "substantially purified" refers to molecules, either nucleic or amino acid sequences, that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated. An "isolated polynucleotide" is therefore a substantially purified polynucleotide.

As used herein, the term "polymerase chain reaction" ("PCR") refers to the method of K.B. Mullis disclosed in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,965,188, all of which are hereby incorporated by reference, which describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing and polymerase extension can be repeated many times (*i.e.*, denaturation, annealing and extension constitute one "cycle"; there can be numerous "cycles") to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" (hereinafter "PCR"). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified."

With PCR, it is possible to amplify a single copy of a specific target sequence in genomic DNA to a level detectable by several different methodologies (*e.g.*, hybridization

with a labeled probe; incorporation of biotinylated primers followed by avidin-enzyme conjugate detection; and/or incorporation of ³²P-labeled deoxynucleotide triphosphates, such as dCTP or dATP, into the amplified segment). In addition to genomic DNA, any oligonucleotide sequence can be amplified with the appropriate set of primer molecules. In particular, the amplified segments created by the PCR process itself are, themselves, efficient templates for subsequent PCR amplifications. Amplified target sequences may be used to obtain segments of DNA (*e.g.*, genes) for the construction of targeting vectors, transgenes, etc.

As used herein, the terms "PCR product" and "amplification product" refer to the resultant mixture of compounds after two or more cycles of the PCR steps of denaturation, annealing and extension are complete. These terms encompass the case where there has been amplification of one or more segments of one or more target sequences.

The term "Central Nervous System" refers to the "spinal cord" and the "brain." The spinal cord comprises white areas and grey areas. The grey area contains nerve cell bodies, whereas the white area is essentially comprised of myelinated nerves. The brain, which is also known as the encephalon, is that portion of the cerebrospinal axis which is contained in the cavity of the cranium. The brain comprises the two cerebral hemispheres, the inter-brain, the mid-brain, the pons Varolli and cerebellum, and the medulla oblongata. The two hemispheres together with the parts derived from the thalamencephalon form the forebrain. The two cerebral hemispheres are separated by the longitudinal fissure and also comprise the Sylvian fissure, the fissure of Rolando, and the parieto-occipital fissure. The lobes on the external surface of the brain comprise the frontal lobe, the parietal lobe, the occipital lobe, and the temporal lobe. Placed along the middle line of the brain are, among others, the rostrum and peduncles of corpus callosum, lamina cinerea, optic commissure and the pituitary body. On each side of the middle line lies the frontal lobe, olfactory lobe and the hemisphere of cerebellum.

The terms "neuron," "neural cell," and "nerve cell" are used interchangeably to refer to a cell which is located in the nervous system. Nerve cells are composed of the nerve cell body (perikaryon), one or more dendrites, and an axon. Neurons can be classified according to the number of processes originating from the cell body. Thus,

unipolar neurons have a single process, bipolar neurons have one axon and one dendrite, while multipolar neurons (which are the most common) comprise more than two processes. The term "neuron" comprises cholinergic neurons and sensory neurons. As used herein, the term "cholinergic neuron" means a neuron in the Central Nervous System (CNS) and in the Peripheral Nervous System (PNS) whose neurotransmitter is acetylcholine. As used
5 herein, the term "sensory neuron" includes a neuron which is responsive to environmental cues (*e.g.*, temperature and movement) from, for example, the skin, muscle and joints of a mammal.

The term "nerve" refers to two or more neurons arranged in linear sequence such
10 that the axon of one neuron establishes a structural and functional link with the dendrite of a second neuron to form a "synapse."

The term "compound" refers to any chemical entity, pharmaceutical, drug, and the like that can be used to treat or prevent a disease, illness, sickness, or disorder of bodily function. Compounds comprise both known and potential therapeutic compounds. A
15 compound can be determined to be therapeutic by testing using the testing methods of the present invention (*i.e.*, a "test compound"). A "known therapeutic compound" refers to a therapeutic compound that has been shown (*e.g.*, through animal trials or prior experience with administration to humans) to be effective in such treatment or prevention. In other words, a known therapeutic compound is not limited to a compound efficacious in the
20 treatment or prevention of neural related disorders.

A compound is said to be "in a form suitable for administration such that the compound is bioavailable in the blood of the animal" when the compound may be administered to an animal by any desired route (*e.g.*, oral, intravenous, subcutaneous, intramuscular, etc.) and the compound or its active metabolites appears in the blood of the
25 animal in an active form. Administration of a compound to a pregnant animal may result in delivery of bioavailable compound to the embryonic progeny of the animal.

The "wild-type β -amyloid" or " β -peptide" gene and gene product refers to the nucleotide and amino acid sequences provided in SEQ ID NOS:6 and 7, respectively. Those skilled in the art will be well aware that certain modifications of SEQ ID NOS:6 and
30 7 can be made which will not interfere with the production of a polypeptide having an

activity indistinguishable from that of the wild-type β -amyloid; the present invention specifically contemplates these variant forms of β -amyloid. A "variant" of the β -peptide is defined as an amino acid sequence that differs by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (*e.g.*, replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes, *e.g.*, replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions (*i.e.*, additions), or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTar software.

The term "an oligonucleotide sequence comprising at least a portion of a human β -amyloid gene" refers to a polynucleotide sequence (*i.e.*, a nucleic acid sequence) containing a nucleotide sequence derived from a human β -amyloid gene. This sequence may encode a portion or all of the β -amyloid protein; alternatively, this sequence may contain non-coding regions derived from the β -amyloid gene or a combination of coding and non-coding regions. The oligonucleotide may be RNA or DNA and may be of genomic or synthetic origin.

As used herein the term "portion" when in reference to a gene refers to fragments of that gene. The fragments may range in size from 10 nucleotides to the entire gene sequence minus one nucleotide. Thus, "an oligonucleotide comprising at least a portion of a gene" may comprise fragments of the gene or the entire gene.

An animal whose genome comprises a "heterologous marker gene" is an animal whose genome contains a marker gene not naturally found in the animal's genome. In one preferred embodiment of the present invention, the heterologous marker gene is a mutant collagen gene, such as the *rol-6(su-1006)* gene. However, it is intended that other marker genes will be used with success in the present invention, including other mutant collagen genes, as well as other marker genes commonly known to those in the art.

As used herein, the term "diagnostic assay" refers to methods for the diagnosis of disease, illness, and/or pathology. It is intended that the term encompass any methods for

diagnosis, including, but not limited to assays based on immunoreactivity (*e.g.*, radioimmunoassays, fluorescence immunoassays, enzyme immunoassays), histochemistry, dye retention or binding (*e.g.*, fixing of dyes such as Congo Red), nucleic acid based diagnostic methods (*e.g.*, identification of nucleic acid sequences associated with disease or pathology), etc.

EXPERIMENTAL

The following examples are provided in order to demonstrate and further illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

In the experimental disclosure which follows, the following abbreviations apply: eq (equivalents); M (Molar); μ M (micromolar); N (Normal); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); $^{\circ}$ C (degrees Centigrade); U (units), mU (milliunits); min. (minutes); sec. (seconds); % (percent); kb (kilobase); bp (base pair); PCR (polymerase chain reaction); β -(1-42) amino acids 1-42 of β -amyloid peptide; Tris (tris(hydroxymethyl)-aminomethane); BSA (bovine serum albumin); Fisher (Fisher Scientific, Pittsburgh, PA); Sigma (Sigma Chemical Co., St. Louis, MO.); Promega (Promega Corp., Madison, WI); Perkin-Elmer (Perkin-Elmer/Applied Biosystems, Foster City, CA); Senetek (Senetek, PLC, Maryland Heights, MO); Boehringer Mannheim (Boehringer Mannheim, Corp., Indianapolis, IN); Stratagene (Stratagene, Inc., La Jolla, CA); and NEB (New England Biolabs, Beverly, MA). Unless otherwise indicated, the restriction enzymes used in these Examples were obtained from NEB. *C. elegans* is available from the Caenorhabditis Genetics Center, at the University of Minnesota, St. Paul, MN.

EXAMPLE 1

Assembly of the β -(1-42) Minigene

1-42) minigene used in subsequent experiments (See, *in C. elegans* engineered to express amyloidic human β -amyloid peptide, as described below, these animals contained constructs in which the 5' promoter/enhancer of *C. elegans* drove the expression of a gene derived from human β -amyloid cDNA clones, as described in *Transgenic Caenorhabditis elegans* (1995), 92:9368-9372 [1995]).

The 42 amino acid β -amyloid peptide derived from human β -amyloid cDNA was assembled. The artificial signal peptide (residues 1-28) was amplified under standard conditions, using "SP-up" (5'-GGCCAAAGGACCC-3') (SEQ ID NO: 1), and "SP-down" (3'-CCAGCAAGAT-3') (SEQ ID NO: 2), cleaved with *NheI* and inserted between the unique *NheI* and *KpnI* sites of construct "pCL2." This process resulted in a recombinant construct such that the signal peptide cleavage site, as predicted by von Heijne, Nucl. Acids Res., 14:4683-4690 [1986], was a Gly-Thr dipeptide encoded by the *KpnI* site.

The 42 amino acid β -amyloid peptide, which contained residues 1-42 of β -amyloid, and which contained a stop codon, was amplified using standard methods, from human β -amyloid cDNA (4T4B (P. Ponte *et al.*, Nature 331:525-527 [1988]), by PCR using primers 5'-GGGGTACCGATGCAGAATTCGACATGA-3') (SEQ ID NO: 3) and 3'-CCCGAGCTCACGCTATGACAACACCGCCAA-3'). The PCR product was cleaved with *KpnI* and *SacI*, and inserted between the unique *KpnI* and *SacI* sites of pCL2, to generate "pCL3."

The 1-42 minigene fragment was removed from this plasmid by PCR and inserted between the unique *NheI* and *SacI* sites of pCL2.

The sequence of the β -(1-42) minigene was confirmed by sequencing the coding strand only, by techniques known in the art. The sequence, SEQ ID NO: 5, is shown in Figure 2. A graphic map of pCL12 is shown in Figure 3. The map shows the DNA and amino acid sequences (SEQ ID NOS: 1-4).

6 and 7, respectively) of pCL12 from nucleotide 1071 through 1253 (*i.e.*, the $\beta(1-42)$ nucleic acid and amino acid sequences) present in the construct.

EXAMPLE 2

Construction of the hsp/GFP Reporter

Example, the expression vector ("pCL25") containing the hsp/GFP reporter expression of the GFP-TT gene was constructed. This vector was used in experiments (See, Example 3) to produce transgenic *C. elegans* was GFP-TT is a modified form of GFP that contains Ser65Thr and Ile167Thr relative to the sequence of the wild-type GFP). The substitutions present in the protein which results in the production of a much brighter than the wild-type GFP.

was constructed as described below. A 431 bp *HindIII*-*Bam*HI fragment of the hsp16-2 promoter was excised from the plasmid "pPD49.78," by digestion with *HindIII*. This 431 bp *HindIII*-*Bam*HI fragment was inserted between the *HindIII* (nucleotide #1) and *Bam*HI (nucleotide #31) sites of the GFP-TT gene in the plasmid "pGFP-TT," (available from Yishi Jin, at the University of California, Santa Barbara). The pGFP-TT plasmid contains the coding regions for GFP-TT inserted into a Tu61 backbone. The DNA sequence of pGFP-TT (SEQ ID NO: 8) is shown in Figure 5. A schematic map of pGFP-TT is shown in Figure 6, in which unique restriction sites are shown. The pGFP-TT plasmid was digested with *HindIII* and *Bam*HI, and the 431 bp fragment containing the hsp16-2 promoter was inserted, in order to construct the hsp-inducible expression vector "pCL25." The DNA sequence of pCL25 and restriction map are shown in Figure 7. Figure 8 provides a schematic map of pCL25 in which unique restriction sites are shown.

EXAMPLE 3

Construction of Transgenic Animals

5 In this Example, the transgenic parent animals were produced, with one line expressing the $\beta(1-42)$ minigene (designated as "CL2005") and the other line expressing the hsp/GFP reporter (designated as "CL2070"). For both lines, the transgenes were introduced into *C. elegans* by gonad microinjection as known in the art and described by Mello *et al.* (Mello *et al.* EMBO J., 10:3959-3970 [1991]). Marker plasmid pRF4 containing the gene (*rol-6*[su1006]) (SEQ ID NO:9) (pRF4 carries a 4 kb *Eco*RI fragment of *C.*
10 *elegans* genomic DNA containing the *rol-6*[su1006] collagen gene in the Bluescribe vector [Stratagene]; See, Mello *et al.*, *supra*; and Kramer *et al.*, Mol. Cell. Biol., 10:2081-2090 [1990]) was coinjected with the constructs, at approximately 100 ng/ μ l for each plasmid, into morphologically wild-type animals, and Roller transgenic progeny were recovered. The (*rol-6*[su1006]) gene (SEQ ID NO:9), contained within the pRF4 plasmid is a mutated
15 *C. elegans* collagen gene, the expression of which produces the dominant, distinctive "Roller" phenotype. Figures 9 and 10 show the DNA and amino acid sequences of the *rol-6* gene used in this Example, respectively.

Transmitting lines were established and maintained by selection for the Roller marker phenotype. Transgenic animals produced in this manner maintain the injected
20 DNA as an extrachromosomal, multicopy array of variable mitotic and meiotic stability. Strains containing chromosomally integrated transgenes were recovered by irradiation of lines containing extrachromosomal transgenic arrays with 7000 rad (1 rad = 0.01 Gy) of gamma rays from a Cesium-66 source. Progeny of irradiated animals were then screen for 100% transmittance of the marker transgene. The transgenes in both the CL2005 and
25 CL2070 lines were chromosomally integrated, and were 100% stable.

Transgenic (*i.e.*, as indicated by the Roller phenotype) animals produce both transgenic (*i.e.*, Roller) and non-transgenic (*i.e.*, non-Roller) progeny. These non-transgenic progeny were found to serve as good internal controls for phenotypic and immunohistochemical comparisons. The expression of GFP in the dual transgenic animals
30 can be detected in less than 24 hours after the upshift of animals from 16 to 25.5°C. at all

stages of development, from late embryonic to adult. Expression of GFP resulted in the production of green fluorescence in muscle cells; intense tissue-specific expression of GFP was observed. The fluorescence can be observed using compound or dissecting epifluorescence microscopy (*i.e.*, with standard fluorescein excitation and emission filters). It is also contemplated that the fluorescence is observable by use of fluorimeters and cell sorters.

EXAMPLE 4

Immunohistochemistry of Transgenic Animals

In this Example, immunohistochemistry was used to confirm the transmittance of the chimeric constructs in large populations of putative integrated lines.

As described by Link (Link, 1995, *supra*), whole mount specimens were prepared by fixing animals in 4% paraformaldehyde and permeabilizing them with 2-mercaptoethanol collagenase as known in the art and described by Link *et al.* (Link *et al.*, Genetics 131:867-881 [1992]). Three antibodies were used (polyclonal rabbit anti- β peptide antibody, Boehringer Mannheim; mouse monoclonal antibody Ab 4G8, available from Senetek; and monoclonal 4.1, a gift from B. Cordell at Scios Nova). The monoclonal 4.1 recognizes residues 8-15 of β -peptide.

As was observed by Link (Link, 1995, *supra*) animals transgenic for the *unc-54*/ β (1-42) minigene construct contained muscle-specific deposits of anti- β peptide immunoreactivity. Non anti- β -immunoreactivity was observed in control animals (*i.e.*, non-transgenic, wild-type *C. elegans* tested concurrently). Although *C. elegans* is reported to contain a homolog of the β -amyloid precursor protein gene (See, Daigle and Li, Proc. Natl. Acad. Sci., 90:2045-2049 [1993]), this sequence does not contain an apparent β -peptide domain. Thus, it would not be expected, nor did it show, cross-reactivity with the anti- β antibody used in these experiments. For dually transgenic animals, immunoreactive deposits accumulated in the body wall of the animals.

Next, to determine whether the immunoreactive deposits observed in the *unc-54*/ β (1-42) strains displayed the tinctural properties of classic insoluble β -amyloid, transgenic

strains were fixed and stained with thioflavin S, a fluorescent amyloid-specific dye, as known in the art, and described by Guntern and Bouras (R. Guntern and C. Bouras, *Experientia* 48:-10 [1992]). Thioflavin S-reactive deposits were found in all strains containing the *unc-54* β -(1-42) minigene constructs, but not in control wild-type animals.

5 Whole mounts of fixed dual transgenic animals stained with coumarin-phalloidin (a muscle-specific probe) and anti- β peptide antibody. Muscle-specific β peptide deposits were observed, and showed a qualitative correlation with the level of GFP expression in these muscle cells (*i.e.*, cells that contained more β -peptide deposits were more green than cells with less β -peptide deposits).

10 In addition, Congo Red and Chrysamine G, two dyes known to interact with β -amyloid and have been reported to interfere with its aggregation are tested on the dually transgenic animals. Anti-oxidants (*e.g.*, vitamin E and ascorbate) are also tested. In these experiments, interference with aggregation of β -peptide, or interaction with β -amyloid are observed.

15 All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention
20 as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

CLAIMS

What is claimed is:

- 5 1. A transparent animal comprising:
- (a) a toxic transgene;
- (b) an inducible promotor operably linked to a reporter gene, wherein the
toxic transgene induces the promotor and wherein the expression of the reporter gene is
detectable *in vivo*.
- 10 2. The transparent animal of claim 1, wherein the inducible promoter operably
linked to the reporter gene is stably integrated into the genome of the animal.
3. The transparent animal of claim 2, wherein the inducible promoter is
15 induced by disruption of cellular function.
4. The transparent animal of claim 2, wherein the inducible promoter is a heat
shock promoter.
- 20 5. The transparent animal of claim 2, wherein the toxic transgene is β -peptide.
6. The transparent animal of claim 2, wherein said transparent animal is
selected from the class Nematoda.
- 25 7. The transparent animal of Claim 6, wherein said transparent animal is
Caenorhabditis elegans.

8. The transparent animal of Claim 2, wherein said genome comprises SEQ ID NO:5 and SEQ ID NO:8.

5 9. The transparent animal of Claim 2, wherein said reporter gene is green fluorescence protein (GFP).

10. The transparent animal of claim 1, further comprising a heterologous gene marker.

10 11. A method for producing a dually transgenic non-human animal comprising:
a) providing:
i) a first and second non-human animal; and
ii) a first toxic transgene; and
iii) a second transgene, comprising a reporter;
15 b) introducing said first toxic transgene into the genome of said first non-human animal to produce a first transgenic animal, and introducing said second transgene into the genome of said second non-human animal to produce a second transgenic animal; and
c) mating said first transgenic animal with said second transgenic animal
20 to produce a dually transgenic animal, wherein said toxic transgene and said reporter are expressed.

12. A dually transgenic non-human animal produced according to the method of claim 11, wherein the toxic transgene is β -peptide.

25

13. A dually transgenic non-human animal produced according to the method of claim 11, wherein said non-human animal is transparent.

14. The dually transgenic non-human animal of claim 13, wherein said non-human animal is a nematode.

5 15 The dually transgenic non-human animal of claim 14, wherein said nematode is *Caenorhabditis elegans*

16. The method of claim 11, wherein said reporter is green fluorescent protein.

10 17. The dually transgenic non-human animal produced according to the method of claim 11, wherein said first transgene comprises pCL25.

18. The dually transgenic non-human animal produced according to the method of claim 11, wherein said second transgene comprises pCL12.

15 19. The dually transgenic non-human animal produced according to the method of claim 11, further comprising a heterologous marker gene.

20. A method for testing compounds for toxicity, comprising:

20 a) providing:

i) a dually transgenic non-human animal expressing a toxic transgene and reporter gene operably linked to a promoter inducible by the toxic transgene;

25 ii) a composition comprising a test compound in a form suitable for administration such that said compound is bioavailable in the cells of said non-human animal; and

b) administering said test compound to said non-human animal.

21. The method according to claim 20, wherein the toxic transgene is β -peptide and the reporter gene is GFP.

5

22. The method of Claim 21, further comprising c) measuring a reduction in the fluorescence of said non-human animal and thereby identifying a compound as therapeutic.

23. The method of Claim 20, wherein said compounds inactivate said β -peptide
10 expressed by said dually transgenic animal.

1/68





Fig. 1

2/68



3/68




Fig. 2A

4/68




Fig. 2B

5/68




Fig. 2C




Fig. 2D

7/68

Sau96 I
Nla IV
Hae III
Ava I
Sec I
Mnl I
SfaN I
CTCATCAGGATCA
GAGTAGCGCTAGT
971
981
981
982
990
990
990
991
997
997
998
998
998
998
Hae I
Hae I
Eae I
Bcl I
Ava II
PpuM I
EcoO109 I
-SP-up
Taq I
BstB I
Xmn I
1014
1015
1038
1040

Sty I
Sec I
Sau96 I
Nla IV
Ava I
PpuM I
EcoO109 I
Mnl I
Mae I
Nhe I
Nsi I
CATTTCAGGAGGACCCCTTGGCTAGC
GTAAAAGTCCTCGGAAACCGGATCGT
1050
1051
1051
1052
1052
1056
1056
1061
1062
1071
1071
Nla IV
Ban I
BspM I
Rsa I
Nla IV
Kpn I
Ban I
Asp718
1109
1109
1116
1119
1119
1119
1119
1120

SfaN I
EcoR I
Nla III
Dde I
Ple I
Hinf I

Mbo II

Fig. 2E

8 / 68

Fig. 2F




Fig. 2G




Fig. 21

13/68



Fig. 2K

14/68





Fig. 2L






Fig. 2N

18/68




Fig. 2P

19/68




Fig. 2Q

20/68




Fig. 2R

22/68




Fig. 2T

23/68




Fig. 3

24/68

1071 /	1	1101 /	11
ATG CAT AAG GTT TTG CTG GCA CTG TTC TTT ATC TTT CTG GCA CCA GCA GGT ACC GAT GCA			
met his lys val leu leu ala leu phe phe ile phe leu ala pro ala gly thr asp ala			
1131 /	21	1161 /	31
GAA TTC CGA CAT GAC TCA GGA TAT GAA GTT CAT CAT CAA AAA TTG GTG TTC TTT GCA GAA			
glu phe arg his asp ser gly tyr glu val his his gln lys leu val phe phe ala glu			
1191 /	41	1221 /	51
GAT GTG GGT TCA AAC AAA GGT GCA ATC ATT GGA CTC ATG GTG GGC GGT GTT GTC ATA GCG			
asp val gly ser asn lys gly ala ile ile gly leu met val gly gly val val ile ala			
1251 /	61		
TGA			
OPA			

Fig. 4

25/68

Fig. 5A
Fig. 5B
Fig. 5C
Fig. 5D
Fig. 5E
Fig. 5F
Fig. 5G
Fig. 5H
Fig. 5I
Fig. 5J
Fig. 5K
Fig. 5L
Fig. 5M
Fig. 5N
Fig. 5O
Fig. 5P
Fig. 5Q
Fig. 5R

Fig. 5

28 / 68






Fig. 5C

29 / 68








Fig. 5G




Fig. 51

Fig. 5J

36 / 68




Fig. 5K




Fig. 5L





Fig. 5M



3942 3971 3976 3992 3993

Scrf I
EcoR II
BstN I
Sec I

4076
4077
4077
4077

4160

4153

4162

4173

4099
4099

4035
4035

4045
4046

4058

4071
4071

4080

4162

4173

4180

4190

4200

4210

4220

4230

4240

4250

4260

4270

4280

4290

4300

4310

4320

4330

4340

4350

4360

4370

4380

4390

4400

4410

4420

4430

4440

4450

4460

4470

4480

4490

4500

4510

4520

4530

4540

4550

4560

4570

4580

4590

4600

4610

4620

4630

4640

4650

4660

4670

4680

4690

4700

4710

4720

4730

4740

4750

4760

4770

4780

4790

4800

4810

4820

4830

4840

4850

4860

4870

4880

4890

4900

4910

4920

4930

4940

4950

4960

4970

4980

4990

5000

5010

5020

5030

5040

5050

5060

5070

5080

5090

5100

5110

5120

5130

5140

5150

5160

5170

5180

5190

5200

5210

5220

5230

5240

5250

5260

5270

5280

5290

5300

5310

5320

5330

5340

5350

5360

5370

5380

5390

5400

5410

5420

5430

5440

5450

5460

5470

5480

5490

5500

5510

5520

5530

5540

5550

5560

5570

5580

5590

5600

5610

5620

5630

5640

5650

5660

5670

5680

5690

5700

5710

5720

5730

5740

5750

5760

5770

5780

5790

5800

5810

5820

5830

5840

5850

5860

5870

5880

5890

5900

5910

5920

5930

5940

5950

5960

5970

5980

5990

6000

6010

6020

6030

6040

6050

6060

6070

6080

6090

6100

6110

6120

6130

6140

6150

6160

6170

6180

6190

6200

6210

6220

6230

6240

6250

6260

6270

6280

6290

6300

6310

6320

6330

6340

6350

6360

6370

6380

6390

6400

6410

6420

6430

6440

6450

6460

6470

6480

6490

6500

6510

6520

6530

6540

6550

6560

6570

6580

6590

6600

6610

6620

6630

6640

6650

6660

6670

6680

6690

6700

6710

6720

6730

6740

6750

6760

6770

6780

6790

6800

6810

6820

6830

6840

6850

6860

6870

6880

6890

6900

6910

6920

6930

6940

6950

6960

6970

6980

6990

7000

7010

7020

7030

7040

7050

7060

7070

7080

7090

7100

7110

7120

7130

7140

7150

7160

7170

7180

7190

7200

7210

7220

7230

7240

7250

7260

7270

7280

7290

7300

7310

7320

7330

7340

7350

7360

7370

7380

7390

7400

7410

7420

7430

7440

7450

7460

7470

7480

7490

7500

7510

7520

7530

7540

7550

7560

7570

7580

7590

7600

7610

7620

7630

7640

7650

7660

7670

7680

7690

7700

7710

7720

7730

7740

7750

7760

7770

7780

7790

7800

7810

7820

7830

7840

7850

7860

7870

7880

7890

7900

7910

7920

7930

7940

7950

7960

7970

7980

7990

8000

8010

8020

8030

8040

8050

8060

8070

8080

8090

8100

8110

8120

8130

8140

8150

8160

8170

8180

8190

8200

8210

8220

8230

8240

8250

8260

8270

8280

8290

8300

8310

8320

8330

8340

8350

8360

8370

8380

8390

8400

8410

8420

8430

8440

8450

8460

8470

8480

849

Fig. 5R

44/68




Fig. 6

1942 Afl II

45/68

Fig. 7A
Fig. 7B
Fig. 7C
Fig. 7D
Fig. 7E
Fig. 7F
Fig. 7G
Fig. 7H
Fig. 7I
Fig. 7J
Fig. 7K
Fig. 7L
Fig. 7M
Fig. 7N
Fig. 7O
Fig. 7P
Fig. 7Q
Fig. 7R
Fig. 7S
Fig. 7T


Fig. 7

46/68


1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 10

Fig. 7A

48/68



49/68



50/68




Fig. 7E

51/68




Fig. 7F

52/68

1208
1209
1210
1210

1273
1275
1276

1360

1362
1376
1383
1390
1390
1391

1407
1407

1446
1451

1477
1481
1482

1507
1519

1520

1575
1576

1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600

1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700

1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800

1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900


1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000

2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100

2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200

2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224

Fig. 7G



55/68




Fig. 7J

58/68

Fig. 7M





Fig. 20






Fig. 79

63/68




Fig. 7R

65/68




Fig. 7T

66/68




Fig. 8

67/68

1 atcaaatga catccttatg tatatgcatt ctttttatgg ttctctgata ttattctct
 61 ctttatacatt tcacttttcc ttggcaactcg tcaaaaaate tcaattaaaa atttctggat
 121 atatcttttag atctaactga aaatttccag atgaccttaa ctacggcgac gtccggcgcc
 181 attgtatttt ctggagccac ttgttggtt tctcttttg cegetgcttc gctttacagt
 241 caagtttcca atatctggaa tgaactggat gcggaatcg ccaacttcag agtgaattc
 301 ctatatagca ttctaatc actgataatt tcaattattt agagtctcac tgaagacatg
 361 tgggttgata tggtaaaact tggagcagga accgcttcca accgtgtgag acgtcaacaa
 421 tatggaggat atggagccac tgggtgtcag ccaccagcac caactccaaa cccatatgga
 481 ggatfatggag caagccagcc agctccacca gagaattcc cagatggtat accaatgga
 541 ggaatcaac caagttccc aggaggtgga ttcccagatg gtccattccc gaatggagga
 601 ggaccacgtg gaggaatca atgtcaatgc actgttgaga actcatgecc accaggacca
 661 gctggaccag aaggagagga aggaccagat gggcacgatg gacaagacgg agtcccagga
 721 ttggacgga aagatgccga agatgttcaa aacacccccc caacaggatg ctcaactgt
 781 ccacaaggac cacttggacc acaaggacca aatggagctc caggactcag aggaatgcgc
 841 ggagctcgtg gacaaccagg acgtccagga agagacggaa accaggat gccaggagac
 901 tgtggaccac caggggcccc aggatecgtat ggaagaccag gatctccagg aggaagggga
 961 gacgatggag agagaccatt gggccgcccc ggaccaagag gaccaccagg agaggctggg
 1021 ccagaaggac cacagggacc aactggaaga gatgttacc caggacagtc tggaccacaa
 1081 ggagagccag gccttcaagg atatggagga gctgctggag aggaacggga ggaatgaaaaa
 1141 aatatataac gacttcttgt aatatataaa atttcagacc agaaggacca ccaggagccc
 1201 caggacttcc aggaagaagt gccgaatact gcaaatgccc aggaagagaa ggagatgctg
 1261 gacgaagtgc cagacgtcat cgcaattcc aattgtagac aaattcatga cattttcca
 1321 aaaaataaaa catactctc aaaattttt gtttgtatt tggttacca tggatgttaa
 1381 gaactttctt ggggaaata attaaatac aaaactgtat a

Fig. 9

68/68

MTLTTATSGAIVFSGATLLVSLFAAASLYSQVSNIWNELDAEIANFRSLTEDMWVDMVK
LGAGTASNRVRRQQYGGYGATGVQPPAPTNPYGGYGASQAPPEKFPDGIPNGGNQP
KFPGGGFDPFPNNGGPRGGNQCQCTVENSPPGAGPEGEGPDGHDGQDGVPGFD
GKDAEDVQNTPTGCFTCPQQPLGPQGNAPGLRGMRGARGQCPGRDGNPGMPG
DCGPPGAPGSDGKPGSPGGKDDGERPLCRPGRGPPGEAGPEGQGTGRDAYPGQSG
PQGEPLQGYGGAAGEDGPEGPPGAPGLPGKDAEYCKCPGREGDAGRSARRHRKFQL

Fig. 10




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶: C12N 15/00, A01K 67/033, C07K 14/435, 14/47</p>	<p>A3</p>	<p>(11) International Publication Number: WO 98/28971</p> <p>(43) International Publication Date: 9 July 1998 (09.07.98)</p>
<p>(21) International Application Number: PCT/US97/23819</p> <p>(22) International Filing Date: 31 December 1997 (31.12.97)</p> <p>(30) Priority Data: 08/778,645 3 January 1997 (03.01.97) US 60/080,222 3 January 1997 (03.01.97) US</p> <p>(71) Applicant (for all designated States except US): UNIVERSITY TECHNOLOGY CORPORATION [US/US]; Suite 250, 3101 Iris Avenue, Boulder, CO 80301 (US).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): LINK, Christopher [US/US]; 1837 Spruce Street, Boulder, CO 80302 (US).</p> <p>(74) Agents: MONROY, Gladys, H. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p> <p>(88) Date of publication of the international search report: 29 October 1998 (29.10.98)</p>

(54) Title: TRANSGENIC ANIMALS FOR *IN VIVO* VISUALIZATION OF BETA-AMYLOID TOXICITY

(57) Abstract

The present invention relates to methods and compositions for visualization of the toxic effects of transgenes *in vivo*. In particular, the present invention provides methods and compositions for the production and use of transgenic, including dually transgenic, *Cuenorhabditis elegans* for visualization of the toxic effects of β -amyloid accumulation *in vivo*.



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 97/23819

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/00 A01K67/033 C07K14/435 C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim(s)
X	CA 2 088 379 A (CANDIDO EDWARD P M : STRINGHAM EVE G (CA); JONES DONALD (CA)) 30 July 1994	1-4, 6, 7, 13-15, 20
Y	see the whole document	1-23
Y	LINK, C.D.: "Expression of human beta-amyloid peptide in transgenic Caenorhabditis elegans" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 92, no. 20, 26 September 1995, pages 9368-9372, XP002065639 WASHINGTON US cited in the application	1-8, 10-15, 18-20, 22, 23
X	see the whole document	11-15
	--- -/-	



Further documents are listed in the continuation of box C



Patent family members are listed in annex

Special categories of cited documents

- A: document defining the general state of the art which is not considered to be of particular relevance
- E: earlier document but published on or after the international filing date
- L: document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- O: document referring to an oral disclosure, use, exhibition or other means
- P: document published prior to the international filing date but later than the priority date claimed

- T: later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- X: document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- Y: document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- S: document member of the same patent family

Date of the actual completion of the international search

2 September 1998

Date of mailing of the international search report

11 09 1998

Name and mailing address of the ISA

European Patent Office P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040 Tx 31 651 eponi
Fax (+31-70) 340-3016

Authorized officer

Chambonnet, F

INTERNATIONAL SEARCH REPORT

Int. Patent Application No

PCT/US 97/23819

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>LINK, C.D. "Transgenic Caenorhabditis elegans as model system to study amyloid formation and toxicity"</p> <p>NEUROBIOLOGY OF AGING.</p> <p>vol. 15, no. suppl. 1, 29 July 1994, page S33.XP002065640</p> <p>see the whole document</p> <p>---</p>	1.5-7
X	<p>WO 95 11968 A (ATHENA NEUROSCIENCES INC ;LILLY CO ELI (US)) 4 May 1995</p> <p>see page 38, line 3 - page 39, line 22</p> <p>---</p>	11.12.19
Y	<p>WO 93 23533 A (TSI CORP) 25 November 1993</p> <p>see page 27, paragraph 3 - page 28, paragraph 2</p> <p>---</p>	11-15
Y	<p>CHALFIE M ET AL: "GREEN FLUORESCENT PROTEIN AS A MARKER FOR GENE EXPRESSION"</p> <p>SCIENCE.</p> <p>vol. 263, no. 5148, 11 February 1994, pages 802-805. XP002003599</p> <p>see the whole document</p> <p>---</p>	1.2.6.7. 9.11. 13-17. 20.21
Y	<p>MOSS, J. B. ET AL. "Green fluorescent protein marks skeletal muscle in murine cell lines and zebrafish"</p> <p>GENE.</p> <p>vol. 173, no. 1, 1996, pages 89-98.</p> <p>XP002065641</p> <p>AMSTERDAM NL</p> <p>see the whole document</p> <p>-----</p>	1.2.9. 13.16.17

INTERNATIONAL SEARCH REPORT

Int'l application No
PCT/US 97/23819

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons

- 1 ☐ Claims Nos
because they relate to subject matter not required to be searched by this Authority, namely
- 2 ☐ Claims Nos
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically
- 3 ☐ Claims Nos
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6 4(a)

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows

see additional sheet

- 1 ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims
- 2 ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee
- 3 ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos
- 4 ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims, it is covered by claims Nos

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest
- ☒ No protest accompanied the payment of additional search fees

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-10 and partially 11-23

a transparent animal comprising :

- (a) a toxic transgene;
 - (b) an inducible promotor operably linked to a reporter gene; wherein the toxic transgene induces the promotor and wherein the expression of the reporter gene is detectable in vivo; and
- a method for providing such a dually transgenic transparent non-human animal comprising :

- a) providing :
 - i) a first and a second non-human transparent animals;
 - ii) a first toxic transgene; and
 - iii) a second transgene, comprising an inducible promotor operably linked to a reporter gene, wherein the toxic transgene induces the promotor and wherein the expression of the reporter gene is detectable in vivo,

- b) introducing said first toxic transgene into the genome of said first non-human animal to produce a first transgenic animal and introducing said second transgene into the genome of said second non-human animal to produce a second transgenic animal; and

- c) mating said first transgenic animal with said second transgenic animal to produce a dually transgenic transparent non-human animal wherein said toxic transgene is expressed and induces the promotor so that the said reporter gene is also expressed and detectable in vivo;

2. Claims: partially 11-23

a method to produce a dually transgenic non-human animal with a toxic transgene and a reporter are expressed comprising :

- a) providing :
 - i) a first and a second non-human animal;
 - ii) a first toxic transgene; and
 - iii) a second transgene, comprising a reporter,

- b) introducing said first toxic transgene into the genome of said first non-human animal to produce a first transgenic animal and introducing said second transgene into the genome of said second non-human animal to produce a second transgenic animal; and

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

c) mating said first transgenic animal with said second transgenic animal to produce a dually transgenic non-human animal wherein said toxic transgene and said reporter are expressed

in the cases non relevant of the first described subject, that means where it is not a transparent animal comprising :

- (a) a toxic transgene;
- (b) an inducible promotor operably linked to a reporter gene; wherein the toxic transgene induces the promotor and wherein the expression of the reporter gene is detectable in vivo.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/23819

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
CA 2088379 A	30-07-1994	NONE	
WO 9511968 A	04-05-1995	US 5604102 A	18-02-1997
		US 5612486 A	18-03-1997
		AU 8079894 A	22-05-1995
		AU 8080994 A	22-05-1995
		CA 2174429 A	04-05-1995
		CA 2174632 A	04-05-1995
		EP 0736106 A	09-10-1996
		EP 0730643 A	11-09-1996
		JP 9508196 T	19-08-1997
		JP 9507746 T	12-08-1997
		WO 9511994 A	04-05-1995
WO 9323533 A	25-11-1993	AU 4240293 A	13-12-1993